

Ettinger & Feldman – Textbook of Veterinary Internal Medicine

Client Information Sheet

Disk Disease

Richard A. LeCouteur

What is disk disease?

The spinal column is made up of a number of small bones called *vertebrae* that are lined up like building blocks. A hole in the center of each vertebra forms a tunnel in which the spinal cord lies. The spinal cord is extremely important because it carries the messages from the brain to the rest of the body. The spinal cord is extremely delicate, and being surrounded by the bony vertebrae helps to protect it. Between each pair of vertebrae, just underneath the spinal cord, is a little cushion, called an *intervertebral disk*. Disks cushion the vertebrae from one another and provide flexibility to the spine during movement.

As a part of the normal aging process, these disks deteriorate, resulting in so-called *disk disease*. Normally, each disk consists of an outer fibrous ring and an inner gelatinous center (a good analogy would be a jelly donut). With age this “doughnut” changes in consistency; the outer fibrous ring becomes fragmented and the inner “jelly” center hardens to a consistency of hard cheese. The fragmented outer fibrous ring may no longer be able to hold this hard center in place, and movement of the vertebrae on either side may suddenly squeeze the disk out of its normal position. Unfortunately, this material usually moves upward and comes to rest against the spinal cord, bruising it in the process.

This “slipping” of the disk often occurs explosively, causing significant damage to the spinal cord and pain to the animal. In this abnormal position the disk presses against the spinal cord, causing further damage.

This type of disk disease may occur in dogs and cats of any age or breed but most commonly in the “short-legged” breeds, such as the following:

- Dachshund
- French bulldog
- Welsh corgi
- Pekingese

What are the symptoms of disk disease?

The parts of the spine most commonly affected by “slipped” disks are the neck and middle to lower back, as the disk “slips” out of place and pushes against the spinal cord.

Some common symptoms of disk disease include the following:

- Significant back pain
- Difficulty in using front and/or rear legs
- Lameness
- Poor coordination
- Paralysis

What tests are needed?

These symptoms indicate that the dog or cat has a problem affecting the spinal cord but not the exact location or cause of the problem. Disk disease, a tumor of the spine, or an infection of the spine may all produce similar symptoms. Tests are needed to determine the exact location and cause of the problem and to decide on the appropriate therapy. To accomplish this, the patient must be anesthetized for x-rays and collection of fluid from around the spinal cord. *Myelography* is an x-ray study in which a special dye is injected into the fluid surrounding the spinal cord. This then allows any disk material pushing against the cord to be identified on the x-rays. Analysis of the fluid around the spinal cord helps to rule out other causes of the problems such as infection.

What treatment is needed?

In most cases disk disease is a problem requiring surgery to remove the disk material compressing the spinal cord. The surgery used most frequently to remove disk material from around the spine is called *laminectomy*. Occasionally, animals with disk disease are not treated by means of surgery. In these animals, *strict cage confinement* and immobilization are used. Usually this approach is used for a first bout of back pain in animals that do not have problems walking. Although strict cage confinement does not correct the spinal cord compression, it may temporarily reduce some of the pain and swelling around the spinal cord and permit the ruptured disk to “heal.”

As time goes on, it is uncommon for animals treated without surgery to suffer repeated bouts of pain, lameness, and paralysis as additional disk material slips and compresses the spinal cord. With each bout of disk disease the spinal cord suffers additional *permanent* damage. Surgical removal of disk material from the spinal canal is the only treatment that provides rapid and maximal recovery of spinal cord function.

Cortisone administration to animals with disk disease is of therapeutic value only during the first 8 hours after the initial spinal cord injury. Current scientific evidence does not support the use of cortisone beyond this time. Furthermore, the adverse effects of cortisone (e.g., stomach ulcers) must always be kept in mind.

What is the prognosis?

For animals undergoing a laminectomy (surgery), the speed of recovery and the extent to which normal function of the legs is regained depend on many factors, including the degree of the damage to the spinal cord and the length of time that the spinal cord has been compressed by the disk material. Animals exhibiting severe neurologic signs (e.g., depressed feeling in their toes), a

rapid onset of symptoms (hours), and a long period of time before surgery generally have a prolonged recovery period and may have varying degrees of permanent damage.

Contacts for Further Information



**Animal Health
Care Center**
of Hershey

**948 E. Chocolate Ave
Hershey, PA 17033
(717) 533-6745**